

## Glucose Monitoring and Insulin Administration Test

Name	e:	Date:	
1.	Which	of the following ARE symptoms of HYPERglycemia? Select all that apply.	
		Increased urine output	
		Cold/ clammy	
		HIGH pulse	
	d.	Increased thirst	
	e.	Headache	
2.	Which	of the following ARE symptoms of HYPOlycemia? Select all that apply.	
	a.	Shakiness	
	b.	Confusion	
	c.	Hunger	
	d.	Cold/Clammy	
	e.	Increased energy	
3.	Insulin	n is measured in:	
	a.	mL	
	b.	Units	
	c.	MEQ	
	d.	Grams	
4.	What	What are the most common subcutaneous insulin injection sites? Select all that apply.	
	a.	Through IV	
	b.	Abdomen	
	c.	Back of upper arm	
	d.	Thigh	
5.	When	giving insulin injections you should ALWAYS: select all that apply.	
	a.	Take the resident to a private area.	
	b.	Cleanse the area with alcohol wipe in a circular motion starting from the outside to the	
		inside.	
	C.	Rotate injection sites	
	d.	Reuse needle	
6.	If a dia	f a diabetic resident complains of feeling weak, seems confused and skin is noted to be	
	pale/clammy, you should:		
	a.	Take them for a walk	
	b.	Ask them to lie down and check on them later	
	c.	Check blood glucose, if low and able to swallow, give juice or candy. If they have glucose ge	
		or lozenges, call the Nurse to get permission to administer.	
	d.	Give insulin	
7.	What	equipment/supplies are needed to assist with administering Insulin?	
	a.	Just give the insulin	
	b.	Gloves, alcohol wipes, MAR, Insulin Pen, Needles and rotating injection site chart	

d. Don't worry, you can gather what you need after you get the resident.

c. Insulin Pen, Needle, MAR

8. When preparing the site for injection, you should?

- a. Give the intended site a swipe with an alcohol pad. b. Just inject no prep needed c. Using an alcohol pad, going in a circular motion, clean from outside to inside area. d. Using an alcohol pad, going in a circular motion, clean from inside to outside area.
- 9. A diabetics diet should consist of:
  - a. 3-4 large meals a day
  - b. Have equal amounts of carbs, sugars, and protein
  - c. Eat whatever and whenever because they are on insulin
  - d. No concentrated sugars, low carbs, low fat, and 5-6 small meals a day
- 10. When using an insulin pen, what is considered best practice to ensure your needle on the pen is working?
  - a. Prime the insulin pen
  - b. Administer the injection and ask the patient if they felt it.
  - c. Do nothing, who cares if the needle is working
  - d. None of the above
- 11. Where do you dispose of the lancet needle after checking the blood glucose level?
- 12. What is the first step in checking a blood sugar?
  - a. Check MAR
  - b. Wash hands
  - c. Gather supplies
  - d. None of the above
- 13. Why is it important to maintain patient privacy during blood glucose check and insulin administration?
- 14. What is the purpose of checking blood sugar BEFORE administering insulin?
  - a. NEVER check blood sugar before insulin administration
  - b. To prevent hypoglycemia (low blood sugar)
  - c. For documenting purposes
  - d. None of the above
  - e. None of the above
- 15. After puncturing the finger for a blood glucose check, you should:
  - a. Wipe the first drop of blood and use the second drop for testing
  - b. Immediately test the initial blood draw
  - c. Puncture again
  - d. None of the above
- 16. How many units of insulin do you prime the insulin pen with?
- 17. You are NOT supposed to clean the glucometer after each use. TRUE or FALSE
- 18. You are supposed to write a open date on insulin pens once used? TRUE or FALSE
- 19. How long is an insulin pen good for once it is opened?
  - a. 45 days
  - b. 30 days
  - c. 14 days
  - d. They do not expire if kept in the fridge
- 20. When cleaning the glucometer, you are to use:
  - a. Alcohol wipes
  - b. Wet wash cloth/paper towel
  - c. Nothing
  - d. Clorox wipes